



Cercus perhaps princeps i Host. but. Missour Left 19. 1870. fust part planeing - flowing wither 5 - 1790 Trunk end speel high, 5 augusted or arager, somewhat astimbeted nanty 4 incheste in diameter, nibs compressed, undéleté cranale. arobe 1/2 mi rigarous shoots 2 inches apart. All die Formula of flower as far as ran be ascertained from the conthered Men man flower Spincher A wary with 25-30 very prominent tuberrules each bearing 2 to mostly 4 Cold bown stout spicing it axil of a mineste soule with however has after Dioppeared . _ One ovary had 10 Hologue riles, Fleente - /10 anaugement - 10 rity on take. The other 8 mon oblige no, 3 amagement, 8 nos on tale lube sinch long, namon, I inch widenry, Color with 8-10 spile bearing arrobae up wands with 6-8 naked ony i axil of mell saly Style thich Stigma rotter away. MISSOURI BOTANICAL copyright reserved cm GARDEN

MISSIMI TOTAMONE GARDEN GEORGE ENGELMANN PAPERS-MISSOURI BOTANICAL copyright reserved cm GARDEN


June 29 1881 Cereus findrefis Shaws Mern. Dol. Gard.

19.54 1870
Without species. Tet. 29 lin /24 Jade 1.30 - ne le or 36 Staur. 39 - 136 Neet 28. 30 Ovany 12 12 21 stongs Thundet areal with brund thought of the spin single (on take som to 2 on 3 spices - quely - axall as fraile smile 12 stiguent erest comment, 9 (. long eco corona








MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS



























copyright reserved cm

BOTANICAL GARDEN




Hort bot Derc: Ze- 1869 1795 Cerent publique. 2, morely 4 willed on Evrhyed ander orrelan with I rule - 5-7 rad.






















Cercus værsable; M. July 19'1959 flower at folie Grieve's Ille flower 7, 21, by, 8/2 inches filly expaled. Lani, Ovag 19-12 l. log, will about 25-30 arrolae, in In arrangement; judville litter lett, lowrrones into I ving short briskle, uppror with 2-3, 2-3 lising Long (- At speins Sept 1818. 4-5 spil sp. with a rette should one Sube 4-5 white les green, smooth, with few Tubundet and low 8 or 10 with 3 - 4 Thing 2-5 C. Co, upprones all of on the d'Cuto park of the spinless protesterances. On the approache the outer ones short redrik green, the imm one feetalood, 9-18 lins by The below 6mg lins, at the face 18 lins in Dameter. Petet i sevent vous, lime lancoly Lithen long 4 livin White, arente, white 40-45 - number. Lowerpart of the tibe verside nakely freble 2 -2/2 inster, monthe, glutstan, hertuniferous; upp part, able I inches tuesely vound with an imme quetity of flames the lower in one, much short thanthe tribe, the outer on, on the face able 1 inch longen The whole forming a beentift druce men, fundstatel extends al intends (see wext sire) flants suite, status gellen obling Hyle Starler at bose graduelly thinner MISSOURI

BOTANICAL copyright reserved cm GARDEN









BOTANICAL copyright reserved GARDEN

MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS

